



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I. If $CKLMN$ be any secant line from C , then $CL \times CM = CK \times CN = CH \times CI$.

II. If $POC'QR$ be any line through C' , secant to both circles, $C'O \times C'R = C'Q \times C'P = C'L \times C'S$.

III. If any circle be drawn to which both circles are either internally or externally tangent, the line through the points of tangency will pass through the external center of similitude, C .

IV. If any circle be drawn to which one of the given circles is internally tangent, and the other externally tangent, the line joining the points of tangency will pass through the internal center of similitude, C' .

Let T be the given point. On the line through CT take a point U such that $CT \times CU = CH \times CI$. This may be done by passing a circle through I , T , and H . It will cut CT in U .

The circle to which both circles are either internally or externally tangent and which passes through T , will also pass through U .

This follows from I. and III.

If through U and T we pass two circles tangent to either of the given circles, (Prob. 17.) they will be tangent to the other.

On the line through $C'T$ take a point V such that $C'T \times C'V = C'S \times C'L$. This may be done by passing a circle through T , L , and S . It will cut $C'T$ in V .

The circle to which one of the given circles is tangent internally and the other externally and which passes through T will also pass through V . This follows from II. and IV.

If through V and T we pass a circle to which one of the given circles is internally tangent, the other will touch it externally and *vice versa*.


Excellent solutions received from *Professors G. B. M. ZERR, P. H. PHILBRICK, and H. C. WHITAKER.*

15. Proposed by ISAAC L. BEVERAGE, Monterey, Virginia.

A man starts from the center of a circular 10 acre field and walks due north a certain distance, then turns and walks south-west till he comes to the circumference, walking altogether 40 rods. How far did he walk before making the turn?

Solution by Professor J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

Let x be the first distance, y the second, and R the radius, then $x + y = 40 \dots (1)$ and $x^2 + y^2 - 2xy \cos 45^\circ = R^2$, or $x^2 - xy\sqrt{2} + y^2 = R^2 = \frac{1600}{\pi} \dots (2)$.



Squaring (1) and subtracting (2) from it, we have $xy=800\left(1-\frac{1}{\pi}\right)$

$(2-\sqrt{2})\dots(3)$. Now we at once get $y-x=40\sqrt{1-2\left(1-\frac{1}{\pi}\right)(2-\sqrt{2})}\dots(4)$.

Combining (1) and (2), we finally obtain

$$x=40\left[1-\sqrt{1-2\left(1-\frac{1}{\pi}\right)(2-\sqrt{2})}\right], \quad y=40\left[1+\sqrt{1-2\left(1-\frac{1}{\pi}\right)(2-\sqrt{2})}\right].$$

Also solved by P. H. PHILBRICK, G. B. M. ZERR, H. M. CASH, P. S. BERG, CHARLES E. MYERS, J. A. CALDERHEAD, SETH PRATT, and H. C. WHITAKER.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

16- Proposed by F. P. MATZ, M. Sc., Ph.D., Professor of Mathematics and Astronomy, in New Windsor College, New Windsor, Maryland.

Differentiate $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$ with regard to $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$.

Solution by J. F. W. SCHEFFER, A. M., Hagerstown, Maryland. and CHARLES E. MYERS, Canton, Ohio.

Let $\frac{2x}{1-x^2}=z$, then $\tan^{-1}\left(\frac{2x}{1-x^2}\right)=\tan^{-1}z$; but $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ by Trigonometry equals $\tan^{-1}\left(\frac{2x}{1-x^2}\right)=\tan^{-1}z$. Hence $\frac{dz}{dz}=1$, that is, since both expressions are identical, the first differential coefficient is=1.

Also solved by Professor MATZ, SCHMITT, and ZERR.

17. Proposed by H. W. DRAUGHON, Clinton, Louisiana.

To find the volume generated by revolving a circular segment whose base is a given chord, about any diameter as an axis.

Solution by the PROPOSER.

In the circle, center C , draw any diameter ECF , and also any chord